

Adiposity and Reproductive Cycling Status in Zoo African Elephants

Daniella E. Chusyd^{1,2}, Janine L. Brown³, Catherine Hambly⁴, Maria S. Johnson¹, Kari Morfeld⁵, Amit Patki⁶, John R. Speakman^{1,2,4,7}, David B. Allison^{1,2,8,9,10}, and Tim R. Nagy^{1,2,9,10}

Objective: The majority of zoo African elephants exhibit abnormal reproductive cycles, but it is unclear why. Acyclicity has been positively associated with body condition scores. The objective of this study was to measure body composition and examine the relationship between adiposity and cyclicity status, mediated by glucose, insulin, leptin, and inflammation.

Methods: Body composition was assessed by deuterium dilution in 22 African elephants. Each elephant was weighed and given deuterated water orally (0.05 mL/kg), and blood was collected from the ear prior to and five times after deuterium administration. Glucose, insulin, leptin, and proinflammatory biomarker concentrations in serum were determined.

Results: Body fat percentage ranged from 5.24% to 15.97%. Fat adjusted for fat free mass (FFM) and age was not significantly associated with cyclicity status ($P = 0.332$). Age was the strongest predictor of cyclicity status ($P = 0.040$). Fat was correlated with weight ($\rho = 0.455$, $P = 0.044$) and when adjusted for FFM with circulating glucose ($\rho = 0.520$, $P = 0.022$) and showed a trend for association with leptin (unadjusted: $\rho = 0.384$, $P = 0.095$; adjusted for FFM: $\rho = 0.403$, $P = 0.087$).

Conclusions: Deuterium dilution appears to be an available technique to measure body composition in African elephants. In this sample, fat was not associated with cyclicity status, and age may be more important to cyclicity status.

Obesity (2018) 26, 103-110. doi:10.1002/oby.22046

Introduction

Obesity is an epidemic not only affecting humans but also animals associated with humans, including companion and domestic animals (1) and, possibly, zoo animals. Approximately 33% of North American female zoo African elephants (*Loxodonta africana*) are classified as obese, based on a body condition score (BCS) (2). BCS is a visual assessment of key skeletal structures and is, therefore, a subjective assessment of subcutaneous body fat stores (3). Although BCS provides an overall assessment of elephant body condition, it does not quantify body composition (fat [FM] or fat free mass [FFM]).

More than 50% of zoo African elephants in the United States exhibit irregular reproductive cycles or are acyclic (4), yet the causes of

acyclicity remain unknown. Previous studies have demonstrated a positive association between condition indices (i.e., BCS and BMI) and rates of reproductive acyclicity in zoo African elephants (5,6), and there is evidence in other species linking body composition to reproductive impairments (7-9). Thus, it is not unreasonable to posit that zoo elephants with increased FM may be more likely to exhibit abnormal reproductive cycles than are elephants with lower FM.

Assessing elephant adiposity is a necessary next step in evaluating this species' reproductive and overall health. Similar reproductive impairments to those observed in zoo elephants have been noted in other species and shown to be associated with excess FM (9,10). The relationship between excess FM and reproductive problems may

¹ Department of Nutrition Sciences, University of Alabama at Birmingham, Birmingham, Alabama, USA. Correspondence: Tim R. Nagy (tnagy@uab.edu)

² Nutrition Obesity Research Center, University of Alabama at Birmingham, Birmingham, Alabama, USA ³ Department of Reproductive Sciences, Conservation & Research Center, National Zoological Park, Smithsonian Institution, Front Royal, Virginia, USA ⁴ Institute of Biological and Environmental Sciences, University of Aberdeen, Aberdeen, Scotland, UK ⁵ Lincoln Children's Zoo, Lincoln, Nebraska, USA ⁶ Department of Biostatistics, University of Alabama at Birmingham, Birmingham, Alabama, USA ⁷ Institute of Genetics and Developmental Biology, Chinese Academy of Sciences, Beijing, China ⁸ Office of Energetics, University of Alabama at Birmingham, Birmingham, Alabama, USA ⁹ Nathan Shock Center, University of Alabama at Birmingham, Birmingham, Alabama, USA ¹⁰ Diabetes Research Center, University of Alabama at Birmingham, Birmingham, Alabama, USA.

Funding agencies: This work was supported in part by the Eppley Foundation for Research, the Nutrition Obesity Research Center (P30DK056336), the Diabetes Research Center (P30DK079626), and the Nathan Shock Center on Aging (P30AG050886). DEC is supported by the National Heart, Lung, and Blood Institute (T32HL105349).

Disclosure: The authors declared no conflict of interest.

Additional Supporting Information may be found in the online version of this article.

Received: 7 August 2017; **Accepted:** 22 September 2017; **Published online** 20 December 2017. doi:10.1002/oby.22046

TABLE 1 Body composition of reproductive-age female zoo African elephants using deuterium dilution by intercept method

ID	Age	BW (kg)	Nd (kg)	TBW (kg)	TBW (%TBM)	FFM (kg)	FM (kg)	WTR (L/d)
A	32	3,436	2,467.22	2,372.32	69.04	3,249.75	186.25	301.01
B	51	4,125	2,874.53	2,763.97	67.00	3,786.26	338.74	260.40
C	30	3,311	2,210.63	2,125.61	64.19	2,911.79	399.21	214.77
D	26	2,850	1,982.87	1,906.61	66.90	2,611.79	238.21	458.38
E	26	3,080	2,178.09	2,094.32	68.00	2,868.93	211.07	317.24
F	26	3,112	2,089.72	2,009.35	64.57	2,752.53	359.47	385.71
G	35	3,833	2,420.62	2,327.52	60.73	3,188.38	644.62	291.87
H	37	4,090	2,942.45	2,829.28	69.17	3,875.73	214.27	441.79
I	43	4,010	2,860.40	2,750.39	68.59	3,767.66	242.34	489.45
L	33	2,979	2,068.54	1,988.99	66.76	2,724.64	254.36	341.28
M	33	3,538	2,263.92	2,176.84	61.53	2,981.98	556.02	234.60
N	37	3,613	2,534.21	2,436.74	67.45	3,337.99	275.01	262.13
P	33	3,973	2,835.94	2,726.87	68.63	3,735.43	237.57	285.99
R	32	3,703	2,612.84	2,512.35	67.85	3,441.57	261.43	338.25
S	34	3,029	2,056.79	1,977.68	65.29	2,709.15	319.85	215.63
T	16	2,211	1,575.02	1,514.44	68.50	2,074.58	136.42	169.45
U	43	4,375	3,020.52	2,904.34	66.38	3,978.55	396.45	319.15
W	48	3,160	2,120.27	2,038.72	64.52	2,792.77	367.23	297.92
X	40	3,920	2,654.58	2,552.48	65.11	3,496.54	423.46	366.13
Y	40	4,465	2,848.31	2,738.76	61.34	3,751.72	713.28	399.08

BW, body weight; Nd, dilution space; TBW, total body water in kilograms and the percent of total body mass; FFM, fat free mass; FM, fat mass; WTR, water turnover rate.

be mediated through the animal's metabolic health. For instance, leptin and insulin are positively associated with FM (11,12). Excess fat is also mechanistically linked with inflammation (13). Leptin, hyperinsulinemia, and chronic inflammation have all been shown to play a role in reproduction and related dysfunction (14,15).

Given the elephant's size, deuterium dilution offers a tenable solution to estimate body composition *in vivo*. Deuterated water has been used to measure total body water (TBW) (i.e., the combination of intra- and extracellular water) in animals ranging in body size from the bumblebee to the Atlantic walrus (16,17). Yet, to our knowledge, no published studies have been conducted in the largest terrestrial mammal, the elephant. Deuterium (^2H), a nonradioactive isotope of hydrogen (^1H), when administered to animals, is diluted by the ^1H in water molecules, providing an estimate of TBW (18). TBW is assumed to be restricted to the animal's FFM compartment; thereby, using the standard mammalian hydration constant, based on TBW of 0.73, FFM can be calculated (19). FM is then inferred by subtracting FFM from weight.

Thus, the purpose of the present study is to measure body composition via deuterium dilution and then to evaluate the relationship between body composition and reproductive cyclicity status in zoo African elephants and to investigate the relationship between body composition, circulating glucose, insulin, and leptin, and inflammatory markers.

Methods

Animals

This study was approved by the Institutional Animal Care and Use Committee of the University of Alabama, Birmingham (UAB) and

by the participating zoos. Of 19 zoos contacted, 13 zoos participated and were visited between November 2014 and June 2016. Data from five of those zoos were omitted because the plateau method for determining TBW was found to be invalid (Supporting Information Table S1). A total of 22 female African elephants of reproductive age (≥ 16 years of age; Table 1) housed in eight accredited US institutions were studied. Elephants were not pregnant. All isotopes were analyzed blind to the animal status.

Body composition

Elephants were weighed to the nearest 5 pounds, pound, or kilogram, depending on each institution's scale. Venous blood was collected from an ear vein by zoo personnel prior to deuterated water administration to determine background isotope enrichment. Subsequently, an oral dose of (99.9% APE) deuterium oxide (0.05 mL D_2O /kg of weight; DLM-4-1000, Cambridge Isotopes, Tewksbury, Massachusetts) was administered using bread (Publix, Birmingham, Alabama) as a vehicle. Bread was weighed (to the closest 0.01 g, Pioneer, Ohaus, Pine Brook, New Jersey), deuterated water was carefully added, and the bread was then reweighed. The difference in weight represented the dose of deuterated water. Each elephant received four to five pieces of bread with approximately 40 to 50 g of deuterated water per piece. Blood (~ 9 mL) was sampled at regular intervals ($\sim 24, 120, 240, 360,$ and 480 hours) post deuterium administration. All samples sat up to 30 minutes in an airtight container to allow for coagulation. Whole blood was centrifuged and serum was collected, aliquoted, and frozen at a minimum of -20°C until shipped on dry ice overnight to UAB. Samples were then kept in a frost-free -80°C freezer until analysis.




Figure 1 Natural log deuterium concentration in venous blood in one reproductive-age female zoo African elephant after enriched orally with deuterated water at Time = 0.

Isotope ratio mass spectroscopy (Finningan Delta V Advantage, Thermo Fisher Scientific, Waltham, Massachusetts) analysis was carried out by the UAB Nutrition Obesity Research Center’s Metabolism Core with guidance and support from the Energetics Research Group at the University of Aberdeen, Aberdeen, Scotland. In brief, the $^2\text{H}/^1\text{H}$ delta value was converted to parts per million and used to calculate deuterium dilution space size (18). The dilution space (N_d) is considered to reflect TBW content, which is then converted to FFM using the hydration constant (see Supporting Information).

To determine blood sampling intervals, in one elephant, blood was collected from an ear vein prior to deuterated water administration and then daily up to 391 hours post administration (Figure 1). These samples were analyzed by liquid water isotope analysis (Los Gatos Research, San Jose, California) at the Energetics Research Group (University of Aberdeen, Scotland).

The water turnover rate was calculated as provided in the Supporting Information.

Determination of reproductive cyclicity status

Reproductive cyclicity status was provided by participating zoos and was based on progestogen analyses of longitudinal serum samples (20). When possible, cycling status was confirmed through longitudinal samples (45% of samples) at the Smithsonian Conservation Biology Institute Endocrinology Laboratory.

BCS

BCS was based on photographs taken as previously described (3). Briefly, photographs of each elephant were taken from three angles (side, rear, and rear-angle view). Two assessors (DEC and KM) visually scored the elephant on a 5-point scale based on key skeletal regions (ribs, pelvic bone, and backbone), where a score of 1 represents the least amount of subcutaneous fat and 5 represents the largest amount of subcutaneous fat (see Supporting Information for details).

Serum analyses

Assays were conducted on the blood samples taken prior to administration of deuterium. The blood was collected in the morning before elephants received their first meal of the day. Elephants’ last meals were given during the late afternoon or evening the day before blood collection. Samples were typically run neat (i.e., not diluted), but if above the assay range, they were diluted in reagent diluent until they fell within the detectable range of the assay.

Serum glucose was measured by an automated glucose analyzer (Stanbio Sirus, Stanbio Laboratories, Boerne, Texas). Samples (3 μL) were done in singlicate and analyzed at the same time.

Serum insulin concentrations were determined with one assay using a solid-phase, two-site bovine insulin enzyme immunoassay (10-1201-01, Mercodia, Uppsala, Sweden) validated for use in African elephants (5). Samples were done in duplicate. The intra-assay coefficient of variation (CV) was 5.2%. See Supporting Information for detailed methods.

Serum leptin concentrations were measured with one assay using a multispecies double-antibody radioimmunoassay (RIA) (XL-85K, Millipore, Billerica, Massachusetts, USA) validated for African elephants (5), which utilized a ^{125}I -human leptin tracer and a guinea pig anti-human leptin antiserum. Samples were done in duplicate. The intra-assay CV was 4.29%. See Supporting Information for detailed methods.

Serum amyloid A (SAA) was determined using an RX Daytona automated clinical chemistry analyzer (Randox Industries-US Ltd., Kearneysville, West Virginia) and commercially available reagents (150 μL) and calibrators (0.1-500 mg/L) with two-level controls (Eiken Chemical Co. Ltd, Tokyo, Japan). Samples (4 μL) were run in singlicate and analyzed at the same time.

Serum tumor necrosis factor alpha (TNF- α) concentrations were measured using an equine TNF- α sandwich enzyme immunoassay kit (ESS0017, Thermo Fisher Scientific; Edwards et al., 2016, unpublished data) according to manufacturer’s instructions. Inter-assay CVs were 6.4% and 2.6% for low and high concentration controls, respectively. CVs for all duplicates were between 0.2% and 13.4%. See Supporting Information for detailed methods.

Statistical analyses

All statistical analyses were performed using SAS 9.4 statistical software (SAS Institute, Cary, North Carolina) and specified prior to examining data, unless otherwise stated. Although 22 elephants were included in this study, 2 elephants were excluded from body composition statistical analyses because blood sample time points were unknown and therefore accurately calculating body composition was not possible.

The primary model to address our main hypothesis was a generalized estimating equation, regressing cyclicity status on FM adjusting for FFM, and age to the power lambda (age^λ). The lambda value for age was calculated by fitting a nonlinear model based on data previously collected on 95 female African elephants, for which age and cyclicity status were known. The best estimate of lambda was 1.62. FM, FFM, and $\text{age}^{1.62}$ were included as continuous variables. To adjust for factors related to residing in the same zoo, the zoo ID

TABLE 2 Sample characteristics of the study sample

	Cycling (n = 13)	Noncycling (n = 9)
Age (y)	31.3 ± 2.1	39.9 ± 1.9 ^a
Body weight (kg)	3,321 ± 137	3,818 ± 176 ^b
Fat mass (kg)	312 ± 34	394 ± 53
Fat free mass (kg)	3,028 ± 145	3,448 ± 173
Height (in)	99 ± 1	101 ± 2
Length (in)	88 ± 2	96 ± 1 ^b
Nulliparous	8/13	7/9
BCS	3.6 ± 0.2	4.3 ± 0.3
Glucose (mg/dL)	78.62 ± 4.52	83.44 ± 2.60
Insulin (µg/L)	0.309 ± 0.043	0.490 ± 0.202
Leptin (ng/mL)	8.53 ± 0.61	8.51 ± 1.12
SAA (mg/L)	0.2 ± 0.1	0.9 ± 0.4
TNF-α (pg/mL)	36.4 ± 9.1	20.6 ± 3.5

Data, except for nulliparous, presented as mean ± SE. Nulliparous data presented as number of elephants that were nulliparous out of total number of elephants.

^aP < 0.01, significance between cycling and noncycling elephants.

^bP < 0.05, significance between cycling and noncycling elephants.

BCS, body condition score; SAA, serum amyloid A; TNF-α, tumor necrosis factor alpha

was treated as a random effect in all the models. The second logistic model added FM² and FFM² to the primary model, as the relationship between body composition and cyclicity status may be nonlinear. Secondary sensitivity analyses were conducted on the primary logistic model after looking at the data. These analyses included the addition of nulliparous status, dominance status, and whether the elephants were housed with male elephants. Nulliparous and dominance status was included as dichotomized variables, while elephants were characterized as either not housed with male elephants, housed with males with direct contact, or housed with males without direct contact.

The mediation of FM in predicting cyclicity status was tested using a two-sided Sobel test (21). Mediators tested were glucose, insulin, SAA, and TNF-α (see Supporting Information).

Pearson correlations between FM and weight, BCS, leptin, glucose, and insulin were conducted. Spearman correlations were conducted between FM and SAA and TNF-α because of the inflammatory biomarkers' non-normal distribution. Correlations between body weight and water turnover, in addition to BCS and weight, FFM, FM, age^{1.62}, and percent body fat, were determined. Partial correlations between FM and leptin, glucose, and insulin, adjusted for FFM, were conducted. Relative fat was determined by the residual for each elephant when FM was regressed on weight.

Descriptive statistics between cycling and noncycling groups were assessed. See Supporting Information for details. Significance was accepted at P < 0.05 (two tailed).

Results

Fifty-nine percent of the elephants exhibited normal reproductive cycles at the time of body composition assessment (n = 13 from eight zoos; mean age 31.3 ± 2.1 years; age range 16-48 years), while the remaining 41% exhibited abnormal reproductive cycles (n = 9 from six zoos; mean age 39.9 ± 1.9 years; age range 33-51 years). Descriptive statistics by cyclicity status are in Table 2.

Body composition and reproductive cyclicity status

Body composition was estimated by deuterium dilution (Table 1) based on the washout curve for each elephant (Supporting Information Figure S1). Body fat percentage averaged 9.69% (SD: 3.18, range: 5.24%-15.97%; n = 20). Generalized estimating equation models analyzing predictors of cyclicity status are in Table 3. FM was not shown to be associated with cyclicity status, either unadjusted (P = 0.131) or adjusted for FFM and age^{1.62} (P = 0.332; Figure 2). When FFM² and FM² were included in the model, FM was not significantly associated with cyclicity status (P = 0.350). When nulliparous status was included, the primary model improved but remained nonsignificant (P = 0.158). The addition of male interaction to the primary model did not improve significance (P = 0.337) unless included with nulliparous status (P = 0.075). When nulliparous and dominance status was included in the primary model, FM was not associated with cyclicity status (P = 0.172). BCS did not

TABLE 3 Estimates for FM in statistical models to predict cycling status

	Estimate	SE	95% CI		P
Cycling = FM	0.005	0.003	-0.002	0.012	0.131
Cycling = FFM FM	0.005	0.004	-0.004	0.014	0.249
Cycling = age ^{1.62} FFM FM	0.004	0.004	-0.004	0.013	0.332
Cycling = age ^{1.62} FFM ² FM ²	0.000	0.000	-0.000	0.000	0.350
Cycling = age ^{1.62} nulliparous FFM FM	0.016	0.011	-0.006	0.037	0.158
Cycling = age ^{1.62} dominant FFM FM	0.004	0.004	-0.004	0.012	0.359
Cycling = age ^{1.62} dominant nulliparous FFM FM	0.014	0.010	-0.006	0.034	0.172
Cycling = age ^{1.62} male FFM FM	0.004	0.004	-0.004	0.013	0.337
Cycling = age ^{1.62} male nulliparous FFM FM	0.046	0.026	-0.005	0.097	0.075

Cycling, cycling status; FFM, fat free mass; FM, fat mass; dominant, dominance status; nulliparous, nulliparous status; male, housed with males with direct contact, housed with males with indirect contact, or not housed with males.




Figure 2 Fat mass, fat mass adjusted by fat free mass, and fat mass adjusted by fat free mass and age by cycling status.

predict cyclicity status either unadjusted ($P = 0.234$) or adjusted for age^{1.62} ($P = 0.750$). Age^{1.62} was a significant predictor of cyclicity status ($P = 0.040$). Weight, unadjusted, was a significant predictor of cyclicity status ($P = 0.038$) but was not significant when adjusted for age^{1.62} ($P = 0.647$).

BCS was strongly correlated with age ($\rho = 0.603$; $P = 0.003$), weight ($\rho = 0.759$; $P < 0.0001$; Figure 3A), FFM ($\rho = 0.702$; $P = 0.001$; Figure 3B), and FM ($\rho = 0.583$; $P = 0.007$; Figure 3C) but not with relative fat ($\rho = 0.256$; $P = 0.276$; Figure 3D) or percent body fat ($\rho = 0.337$; $P = 0.146$).

Body composition and glucose, insulin, leptin, and inflammatory biomarkers

The correlation between FM and relative fat with glucose ($\rho = 0.379$, $P = 0.100$; $\rho = 0.555$, $P = 0.011$, respectively; Figure 4A-4B), insulin ($\rho = 0.369$, $P = 0.110$; $\rho = 0.352$, $P = 0.128$, respectively; Figure 4C-4D), and leptin ($\rho = 0.384$, $P = 0.095$; $\rho = 0.399$, $P = 0.081$, respectively; Figure 4E-4F) nearly reached significance. FM, adjusted for FFM, was correlated with glucose ($\rho = 0.520$; $P = 0.022$) and trended toward significance with insulin ($\rho = 0.371$; $P = 0.118$) and leptin ($\rho = 0.403$; $P = 0.087$). FM was not correlated with SAA ($\rho = 0.007$; $P = 0.979$; Figure 4G) or TNF- α ($\rho = -0.0353$; $P = 0.883$; Figure 4H). Weight was not correlated with water turnover rate ($\rho = 0.357$; $P = 0.123$). Glucose was correlated with insulin ($\rho = 0.430$; $P = 0.046$).

Cyclicity status and glucose, insulin, leptin, and inflammatory biomarkers

Glucose ($P = 0.366$), insulin ($P = 0.406$), leptin ($P = 0.991$), SAA ($P = 0.095$), and TNF- α ($P = 0.349$) did not significantly differ by cyclicity status (Table 1). Although there was no overall association between FM and cyclicity status, mediation analyses were still conducted to better understand why the variables did not relate. Glucose ($P = 0.276$) and insulin ($P = 0.220$) were not mediators between FM and cyclicity status. SAA, analyzed as a dichotomous ($P = 0.564$) and continuous variable ($P = 0.477$), and TNF- α , analyzed as a

dichotomous ($P = 0.841$) and continuous variable ($P = 0.432$), were not mediators between FM and cyclicity status.

Discussion

This study investigated the association between adiposity with reproductive cyclicity status, mediated through glucose, insulin, leptin, and proinflammatory biomarkers. We did not find that FM was significantly associated with cyclicity status in zoo African elephants, nor was the relationship mediated through inflammation, glucose, or insulin. FM was correlated with glucose and nearly correlated with leptin levels. Noncycling elephants were more likely to be older.

Deuterium dilution by the intercept method appears to be a useful, noninvasive technique that can be used to estimate body composition of African elephants. Deuterium dilution measures TBW by using either the plateau or the intercept method (22). The plateau method assumes equilibrium is reached rapidly and neither deuterium nor body water is metabolized during equilibrium (18). These assumptions may be violated in elephants because of their large body volume and slow heart rate (23) (Supporting Information Table S1). The intercept method does not rely on a measurement during equilibrium and therefore allows for a longer equilibrium period and continuous water turnover over an extended period of time through the collection of several samples over time and back extrapolating to when deuterated water was administered (22). The intercept method was therefore used in the present study, in which we observed a log-linear elimination in deuterium enrichment over time in the elephants.

The derivation of FM from TBW estimated by deuterium dilution is based on the ratio of TBW to FFM, termed the hydration constant (19), which to our knowledge has not been determined in elephants. Therefore, we used the most regularly used hydration constant of ~ 0.73 (24). Pace and Rathbun (25) first recommended this hydration constant based on several species of mammals, and since then it has been reinvestigated and confirmed (24). However, the hydration constant may vary by body size. Pitts and Bullard (26) demonstrated that as body size increases, there is a slight decrease in the TBW/FFM. This may be related to larger mammals having a larger proportion of their body mass comprised of bone, as skeletal tissue is comparatively “dry” (26). This could certainly pertain to the African elephant, which may have a hydration constant different from 0.73. For instance, if the hydration constant was 0.71, body fat for this sample would be approximately 3% to 14%. If the hydration constant was 0.75, body fat for this sample would be approximately 8% to 18%. Even though the exact hydration constant of the elephant is unknown, it should not affect the primary results, as any change in the hydration constant would result in a linear transformation of the FM values.

Body fat percentages in this study sample ranged from approximately 5% to 16%. Most African ungulates have little subcutaneous fat, which may help facilitate heat dissipation (27). Most published studies examine only small areas of the body, which show limited quantities of fat, and may be inappropriate to extrapolate to total body fat. For example, female giraffes have been found to have 0.52% to 1.39% rump fat (27). Similarly, the rump fat percentage of male kudus and blesbok is 1.3% and 1.4%, respectively (28).




Figure 3 (A) Relationship between BCS and body weight. (B) Relationship between BCS and FFM. (C) Relationship between BCS and fat mass. (D) Relationship between BCS and relative fat. Relative fat was determined by the residual for each elephant when fat mass was regressed on body weight.

Comparatively more work has been done with domesticated animals. For instance, horses have on average 5.1% to 22.3% body fat, which is correlated with rump fat (29). Elephants have a low surface area to volume ratio, so it would be expected that elephants may have little subcutaneous fat; however, the study sample is from zoos where food is plentiful and of high quality. To determine the accuracy of our body composition measures would require comparing our technique to carcass analysis, which is not feasible.

Obesity has been associated with anovulation in other mammalian species (7,30,31), and in previous studies a relationship between body condition indices (e.g., BMI and BCS) and cyclicity status has been observed in zoo African elephants (5,6). However, the current study did not find such a relationship between adiposity and cyclicity status, which could be because of a variety of factors. For

instance, our sample size ($n = 20$) was smaller than in our previous work (5), which examined 50 elephants (half cycling, half not). Based on our data, to have 0.80 power to detect an association between FM and cyclicity status would require approximately 170 elephants. Differences in results may be age related, which is supported by a recent study demonstrating that age was positively associated with acyclicity (4). We found noncycling elephants to be significantly older and heavier than those that were cycling, with age and weight strongly positively correlated. In our previous study (5), we did not adjust for age, which might explain the discordance in results. Further, BCS correlated most strongly with weight. Therefore, it appears BCS may be capturing the overall weight and not the amount of fat of the elephant. As age and weight are highly correlated ($\rho = 0.679$), age may be driving the relationship between BCS and cyclicity status and should be considered in future studies.




Figure 4 (A) Relationship between fat mass and glucose. (B) Relationship between relative fat mass and glucose. (C) Relationship between fat mass and insulin. (D) Relationship between relative fat mass and insulin. (E) Relationship between fat mass and leptin. (F) Relationship between relative fat mass and leptin. (G) Relationship between fat mass and SAA. (H) Relationship between fat mass and TNF- α . Closed circles: cycling elephants; open circles: noncycling elephants. Relative fat was determined by the residual for each elephant when fat mass was regressed on body weight.

The elephant's metabolic state, rather than absolute FM, may be more important to consider in relation to reproductive and overall health. FM, adjusted for FFM, was positively correlated with

glucose levels, and FM, adjusted and unadjusted, almost reached significance with insulin levels. There was one insulin outlier, and although there was no reason to exclude it, running the analyses without this data point resulted in a significant correlation between unadjusted and adjusted FM and insulin ($\rho = 0.479$, $P = 0.038$; $\rho = 0.516$, $P = 0.028$, respectively). In a postprandial state, elevated blood glucose levels stimulate insulin secretion (32), in turn promoting glucose uptake and utilization and suppressing gluconeogenesis (32); therefore, abnormal insulin secretion, in addition to insulin resistance, can lead to hyperglycemia (32). Hyperglycemia and hyperinsulinemia are both associated with obesity and comorbidities (32,33). Although FM may not be associated with acyclicity, it may impact health in other ways, such as increasing the risk of arthritis and calving problems (34).

Hyperglycemia and hyperinsulinemia stimulate an inflammatory state (35,36), and in other species, inflammation is associated with reproductive impairments and obesity (15). However, our results do not indicate an inflammatory state. The SAA reference interval for clinically healthy Asian elephants is 0 to 47.5 mg/L (37). No elephant, cycling or noncycling, had levels greater than 3.5 mg/L, suggesting all the elephants in this study had low levels of inflammation. Further, SAA levels were not correlated with FM, adjusted or unadjusted, even though SAA is the most sensitive acute phase protein in elephants (37). TNF- α was not associated with FM, adjusted or unadjusted. Taken together, results suggest these elephants did not have chronic inflammation, as observed in other species that exhibit obesity and reproductive impairments (15,38), at least measured by these inflammatory factors; thus, these elephants may not be obese.

FM should be monitored as it is an active endocrine organ. In addition to other adipokines, white adipose tissue produces leptin (39). Leptin plays a permissive role in activating the reproductive axis and, when levels are abnormally high as observed in an obese state (39), prevents ovarian steroidogenesis, inhibiting proper follicle development (40). This is likely not a reason for acyclicity in these elephants, as leptin levels were similar between the two groups; however, as leptin nearly correlated significantly with FM, there may be other related health issues to consider. For example, similar to girls with obesity (39), zoo elephants reach puberty at earlier ages than their wild counterparts (10). Earlier puberty will likely expose the elephants to more reproductive cycles and associated endogenous hormones over their lifetime, which may lead to the development of reproductive tract pathologies (10). With known endocrine function, continued fat accrual may possibly lead to a point at which elephants show similar metabolic dysfunction exhibited by humans and domestic animals with obesity.

This study demonstrated that deuterium dilution can be used to estimate body composition in African elephants. We used a method that has proven accurate over the decades in other species with a 10⁵ times difference in body size. Thus, although assumptions were made (e.g., hydration constant) and there was an inability to validate the technique by total carcass analysis, the method appears to be robust over time and species. Results open up a new avenue of research questioning for large herbivores and the effects of diet and exercise on adiposity and its relation to health and reproduction.

In conclusion, obesity and inflammation were not unequivocally found to be significant factors for this study sample regarding

acyclicity status. Regardless, there does appear to be a relationship between FM and metabolic health in African elephants. The majority of these elephants appear to be metabolically healthy, but some could be categorized as overweight or at risk for metabolic dysfunction. This supports the need to individualize management strategies, as each elephant responds uniquely to the environment and diet. This is of paramount consideration, as the zoo elephant population is currently not self-sustaining. **O**

Acknowledgments

The authors thank Dr. Barbara Gower, Maryellen Williams, Heather Hunter, and Cindy Zeng at the UAB Nutrition Obesity Research Center's Metabolism Core for their assistance with hormone assays and mass spectroscopy and Dr. Katie Edwards at the Smithsonian Conservation Biology Institute for inflammatory analyses. The authors thank Zoo Atlanta, Dallas Zoo, Jacksonville Zoo and Gardens, Maryland Zoo in Baltimore, Memphis Zoo, Zoo Miami, Nashville Zoo at Grassmere, North Carolina Zoo, Pittsburgh Zoo and PPG Aquarium, Riverbanks Zoo and Botanical Garden, San Diego Zoo Global, and Wildlife Safari for their agreement to participate in this study. Specifically, thank you to the zoos' elephant keepers and elephants. They made this study possible. A special thank you to the Birmingham Zoo and Pat Flora and his elephant team for their continuous support, help, and input with method improvement. The opinions expressed herein are those of the authors and not necessarily those of any other organization with which the authors are affiliated.

© 2017 The Obesity Society

References

- Klimentidis YC, Beasley TM, Lin H-Y, et al. Canaries in the coal mine: a cross-species analysis of the plurality of obesity epidemics. *Proc Biol Sci* 2011;278:1626-1632.
- Morfeld KA, Meehan CL, Hogan JN, Brown JL. Assessment of body condition in African (*Loxodonta africana*) and Asian (*Elephas maximus*) elephants in North American zoos and management practices associated with high body condition scores. *PLoS One* 2016;11:e0155146. doi:10.1371/journal.pone.0155146
- Morfeld KA, Lehnhardt J, Alligood C, Bolling J, Brown JL. Development of a body condition scoring index for female African elephants validated by ultrasound measurements of subcutaneous fat. *PLoS One* 2014;9:e93802. doi:10.1371/journal.pone.0093802
- Brown JL, Paris S, Prado-Oviedo NA, et al. Reproductive health assessment of female elephants in North American zoos and association of husbandry practices with reproductive dysfunction in African elephants (*Loxodonta africana*). *PLoS One* 2016;11:e0145673. doi:10.1371/journal.pone.0145673
- Morfeld KA, Brown JL. Ovarian acyclicity in zoo African elephants (*Loxodonta africana*) is associated with high body condition scores and elevated serum insulin and leptin. *Reprod Fertil Dev* 2016;28:640-647.
- Freeman EW, Guagnano G, Olson D, Keele M, Brown JL. Social factors influence ovarian acyclicity in captive African elephants (*Loxodonta africana*). *Zoo Biol* 2009;28:1-15.
- Vick M, Sessions D, Murphy B, Kennedy E, Reedy S, Fitzgerald B. Obesity is associated with altered metabolic and reproductive activity in the mare: effects of metformin on insulin sensitivity and reproductive cyclicity. *Reprod Fertil Dev* 2006;18:609-617.
- Brothers KJ, Wu S, DiVall SA, et al. Rescue of obesity-induced infertility in female mice due to a pituitary-specific knockout of the insulin receptor. *Cell Metab* 2010;12:295-305.
- Ovesen P, Rasmussen S, Kesmodel U. Effect of prepregnancy maternal overweight and obesity on pregnancy outcome. *Obstet Gynecol* 2011;118:305-312.
- Holt WV, Brown JL, Comizzoli P, eds. *Reproductive Sciences in Animal Conservation: Progress and Prospects*. New York, NY: Springer; 2014.
- Krotkiewski M, Seidell J, Björntorp P. Glucose tolerance and hyperinsulinaemia in obese women: role of adipose tissue distribution, muscle fibre characteristics and androgens. *J Intern Med* 1990;228:385-392.
- Considine RV, Sinha MK, Heiman ML, et al. Serum immunoreactive-leptin concentrations in normal-weight and obese humans. *N Engl J Med* 1996;334:292-295.
- Yuan M, Konstantopoulos N, Lee J, et al. Reversal of obesity- and diet-induced insulin resistance with salicylates or targeted disruption of Ikk β . *Science* 2001;293:1673-1677.
- Mitchell M, Armstrong D, Robker R, Norman R. Adipokines: implications for female fertility and obesity. *Reproduction* 2005;130:583-597.
- Robker RL, Wu LL-Y, Yang X. Inflammatory pathways linking obesity and ovarian dysfunction. *J Reprod Immunol* 2011;88:142-148.
- Acquarone M, Born EW. Estimation of water pool size, turnover rate and body composition of free-ranging Atlantic walrus (*Odobenus rosmarus rosmarus*) studied by isotope dilution. *J Mar Biol Assoc U.K.* 2007;87:77-84.
- Wolf T, Ellington C, Davis S, Feltham M. Validation of the doubly labelled water technique for bumblebees *Bombus terrestris* (L.). *J Exp Biol* 1996;199:959-972.
- Speakman JR. *Doubly Labelled Water: Theory and Practice*. New York, NY: Springer; 1997.
- Wang Z-M, Pierson R, Heymsfield SB. The five-level model: a new approach to organizing body-composition research. *Am J Clin Nutr* 1992;56:19-28.
- Brown JL, Walker SL, Moeller T. Comparative endocrinology of cycling and non-cycling Asian (*Elephas maximus*) and African (*Loxodonta africana*) elephants. *Gen Comp Endocrinol* 2004;136:360-370.
- Baron RM, Kenny DA. The moderator-mediator variable distinction in social psychological research: conceptual, strategic, and statistical considerations. *J Pers Soc Psychol* 1986;51:1173. doi:10.1037/0022-3514.51.6.1173
- Speakman JR, ed. *Body Composition Analysis of Animals: A Handbook Of Non-Destructive Methods*. Cambridge, UK: Cambridge University Press; 2001.
- Benedict FG, Lee RC. The heart rate of the elephant. *P Am Philos Soc* 1936;76:335-341.
- Wang Z, Deurenberg P, Wang W, Pietrobelli A, Baumgartner RN, Heymsfield SB. Hydration of fat-free body mass: review and critique of a classic body-composition constant. *Am J Clin Nutr* 1999;69:833-841.
- Pace N, Rathbun EN. Studies on body composition III. The body water and chemically combined nitrogen content in relation to fat content. *J Biol Chem* 1945;158:685-691.
- Pitts G, Bullard T. Some interspecific aspects of body composition in mammals. In: *Body Composition in Animals and Man*. Washington, DC: National Academy of Sciences; 1968:45-70.
- Hall-Martin A, Von la Chevallerie M, Skinner J. Carcass composition of the giraffe *Giraffa camelopardalis giraffa*. *S Afr J Anim Sci* 1977;7:55-64.
- Huntley B. Carcass composition of mature male blesbok and kudu. *S Afr J Anim Sci* 1971;1:125-128.
- Kearns C, McKeever K, Abe T. Overview of horse body composition and muscle architecture: implications for performance. *Vet J* 2002;164:224-234.
- Jungheim ES, Moley KH. Current knowledge of obesity's effects in the pre- and periconceptional periods and avenues for future research. *Am J Obstet Gynecol* 2010;203:525-530.
- Clark A, Ledger W, Galletly C, et al. Weight loss results in significant improvement in pregnancy and ovulation rates in anovulatory obese women. *Hum Reprod* 1995;10:2705-2712.
- Sonksen P, Sonksen J. Insulin: understanding its action in health and disease. *Br J Anaesth* 2000;85:69-79.
- Mehran AE, Templeman NM, Brigidi GS, et al. Hyperinsulinemia drives diet-induced obesity independently of brain insulin production. *Cell Metab* 2012;16:723-737.
- Hermes R, Saragusty J, Schaftenaar W, Göritz F, Schmitt DL, Hildebrandt TB. Obstetrics in elephants. *Theriogenology* 2008;70:131-144.
- Chen J, Wildman RP, Hamm LL, et al. Association between inflammation and insulin resistance in US nondiabetic adults. *Diabetes Care* 2004;27:2960-2965.
- Aronson D. Hyperglycemia and the pathobiology of diabetic complications. In: Fisman EZ, Tenenbaum A, eds. *Cardiovascular Diabetology: Clinical, Metabolic and Inflammatory Facets*. Basel, Switzerland: Karger Publishers; 2008:1-16.
- Isaza R, Wiedner E, Hiser S, Cray C. Reference intervals for acute phase protein and serum protein electrophoresis values in captive Asian elephants (*Elephas maximus*). *J Vet Diagn Invest* 2014;26:616-621.
- Adams AA, Katepalli MP, Kohler K, et al. Effect of body condition, body weight and adiposity on inflammatory cytokine responses in old horses. *Vet Immunol Immunopathol* 2009;127:286-294.
- Blüher S, Mantzoros CS. Leptin in reproduction. *Curr Opin Endocrinol Diabetes Obes* 2007;14:458-464.
- Duggal PS, Van der Hoek KH, Milner CR, et al. The in vivo and in vitro effects of exogenous leptin on ovulation in the rat. *Endocrinology* 2000;141:1971-1976.